

Cataract Removal Discharge Instructions

561-296-2010

Craig A. Skolnick, M.D.

Starting 4 hours after surgery at _____

Lift shield today for drops; remove tomorrow morning. Wear shield at bedtime for 1 week.

VIGAMOX (Tan Top-small bottle) 1 drop 4 times a day

NEVANAC (Gray Top-small bottle) 1 drop 4 times a day

OMNIPRED (Pink Top-milky drop) 1 drop 4 times a day

OTHER: _____

STOP VIGAMOX DROPS AFTER 1 WEEK (OR WHEN FINISHED)

**OMNIPRED AND NEVANAC 4 TIMES A DAY FOR 2 WEEKS THEN
2 TIMES A DAY FOR 2 WEEKS (USE UNTIL BOTTLES ARE EMPTY)**

OMNIPRED, additional NEVANAC sample, paper tape and artificial lens ID card given at surgery center

USE DROPS AT LEAST 5 MINUTES APART FROM EACH OTHER

NO HEAVY LIFTING, SWIMMING OR BENDING YOUR HEAD BELOW THE WAIST FOR 1 WEEK

IT IS NORMAL TO HAVE BLURRED, DIM (for the first 2-3 days the night-time vision may be very dark) OR RED/PINK VISION.

YOU MAY HAVE A HEADACHE, MILD SORENESS OR FOREIGN BODY SENSATION.

YOU ALSO MAY EXPERIENCE FLICKERING OF VISION.